



Curtis Bush, MD
817-878-5300

Proximal Humerus Fracture ORIF

Ultra sling for weeks 0-6

Phase 0 – (QUIET) Days 0-10

Elbow/wrist/hand ROM

Begin active scapular retraction/protraction exercises with therapist cueing

Pendulums to warm-up beginning 7 days post op, incorporate into HEP

Phase 1 -(PASSIVE) Start formal PT 10 days – week 6

Pendulums to warm-up

Supine External Rotation -0° - 30° beginning at 3 weeks with progression to full PROM by 8 weeks

Supine Forward Elevation -0° - 90° beginning at 3 weeks with progression to full PROM by 8 weeks

** progress to upright as tolerated with ER and FE*

Phase 2 -(ACTIVE) Week 7-12

Pendulums to warm-up.

Active Range of Motion with terminal stretch

Supine External Rotation – after 8 weeks progress GRADUALLY to full

Supine Forward Elevation – after 8 weeks; progress GRADUALLY to full

Begin active biceps

Internal Rotation-Full (begin behind the back)

Begin AROM in supine and progress to upright

Phase 3 -(RESISTED) Week 13

Pendulums to warm up and continue with phase 2.

External and Internal Rotation

Standing forward punch

Seated rows

Shoulder shrugs and Biceps curls

Initiation of Interval Sport Programs Surgeon Comments:

Golf: 5-6 months

Tennis: 7-8 months

Ski: 7-8 months